

Stream Quality Monitoring 2015 Annual Report

Kokosing State Scenic River


Contents

Introduction	1
Overview	2
Stream Quality Monitoring Station Map	3
2015 Stream Quality Monitoring Participants.....	4
Station Descriptions	5
Sampling Results and General Trends	7
Total Suspended Solids (TSS).....	8
Comparisons of Collected Stream Quality Monitoring Data.....	9
Table 1 - Macroinvertebrate Pollution Tolerance	9
Tables 2, 3 - 2015 Mean CIVs by Reference Station.....	10
Figure 1.1 - 2015 Mainstem CIV Maximum and Minimum Ranges.....	10
Figure 2.1 - 2006-2015 Mainstem Mean CIVs	11
Figure 1.2 - 2015 North Branch CIV Maximum and Minimum Ranges	12
Figure 2.2 - 2006-2015 North Branch Mean CIVs.....	12
Tables 4, 5 - Qualitative Habitat Evaluation Index (QHEI)	13
Appendix - 2015 Stream Quality Monitoring Data by Monitoring Station	14-15

Introduction

Ohio Scenic Rivers Program

With more than 60,000 miles of streams, Ohio is a water-rich state. Many of Ohio's streams support thriving plant and animal communities, including Ohio's state designated scenic rivers. Administered by the Ohio Division of Watercraft, the Ohio Scenic Rivers Program oversees 14 state designated scenic river systems, comprising 800 river miles along 26 stream segments. These streams represent some of the best of Ohio's waterways.



Stream Quality Monitoring Project

Developed in 1983, the Ohio Stream Quality Monitoring (SQM) Project uses volunteers who assist in aquatic macroinvertebrate monitoring to compile biological and water quality data on the state's scenic rivers. The Ohio SQM Project is an excellent, simple, and cost-effective method of assessing a stream's health.

Aquatic macroinvertebrate organisms lack a backbone (invertebrate), are large enough to view with the naked eye (macro), and spend at least a portion of their lives in the water (aquatic). Macroinvertebrates, such as various aquatic insects (e.g. mayfly, stonefly), are good indicators of stream health. When negative impacts to a stream occur, the result may show a decline or absence of certain macroinvertebrate species.

Through consistent monitoring, changes observed in the macroinvertebrate community help the Ohio Scenic Rivers Program in detecting and addressing potential impacts to a stream. The Ohio Scenic Rivers Program compiles volunteer field assessment information into a statewide database. The database serves as a tool to track short- and long-term changes and trends over time.

SQM Project Relies on Volunteers

Coordinated by the Division of Watercraft's Scenic Rivers Program, the Ohio SQM Project provides opportunities for public participation in scenic river protection efforts. Many local, youth and conservation organizations, individuals, and families are committed to monitoring more than 150 stations along Ohio's scenic rivers.

SQM volunteers collect macroinvertebrate data from selected monitoring stations, also referred to as monitoring sites or reference stations, three times during the monitoring season. Volunteers complete field assessment forms that document taxonomy, tolerance and abundance of collected organisms.

SQM Annual Report

The information collected by volunteers has become a critical tool for documenting the health of Ohio's state scenic, wild, and recreational rivers. This report is a compilation of field data collected during 2015 by volunteers and staff. It also represents a year of dedication and commitment shown to Ohio's special waterways by thousands of SQM volunteers.

Overview

The Kokosing River watershed is situated between two urban centers, Mansfield and Columbus, in the central northeast Ohio rural countryside. Most of the watershed lies within Knox County, which lies on the outer edge of an area once covered by continental glaciers. Two main glacial events left their mark on the landscape. The first was the Illinoian glacier that moved across the watershed and covered most of the eastern part of Knox County. The ice flowed around and between the hills leaving glacial deposits of varying thickness. The second glacial event only covered the western part of Knox County. It also left thick deposits of gravel, sand, and silt.


Forty-eight miles of the Kokosing River system were designated as Ohio's 11th scenic river in 1997. The mainstem Kokosing is designated from the confluence with the Mohican and Walhonding Rivers, totaling 41.1 miles upstream to the Knox/Morrow County line. The North Branch Kokosing is designated from the confluence with the mainstem of the Kokosing River, upstream 6.5 miles, to the confluence with the East Branch Kokosing, just north of Fredericktown. More than 70 species of fish, five species of freshwater mussels, and 24 species of amphibians including the state endangered hellbender inhabit the Kokosing River system. Nearly 90 species of breeding birds also may be found in the forested lands along the riverbanks. These forested banks are a vital part of the Kokosing's exceptional habitat, water quality, and scenery.

The Kokosing River name has an interesting history. The name "Kokosing" in the Delaware Indian language is said to mean "river of many Delaware villages." Other sources say that the word "Kokosing" was used by the Algonquin Indians to mean "river of little owls." In addition, many early historians referred to the river as "Owl Creek." Use of this name prior to 1866 is corroborated by the names used for the Owl Creek Bank of Mt. Vernon, Owl Creek Baptist Church and the Owl Creek Bookstore among others. There are also records of the name "Vernon River" being promoted by Bishop Philander Chase, founder of Kenyon College.

Over two centuries have passed since the first pioneers settled the Kokosing watershed. Remarkably, the population has not quite doubled since 1840. This slow growth rate combined with fairly stable land-use patterns has benefited the river system. Since the initial period of change, the river system has adapted fairly well and has been able to maintain a higher quality than many other Ohio streams. This high quality status may be attributed largely to the diverse stream substrate left by the glaciers and to the presence of a healthy wooded corridor maintained by landowners over the years.

The Kokosing River and North Branch of the Kokosing River are popular streams for canoeing, fishing, bird watching, and other outdoor activities. Additional information about public access facilities on the Kokosing is available by contacting the Division of Watercraft at 740-548-5490 or visiting online at watercraft.ohiodnr.gov .





2015 Stream Quality Monitoring Participants

Whether their contribution was a one-time event or a recurring adventure in stream exploration, the individuals and organizations listed below played a significant role in the preservation of the Kokosing River. Their time and dedication to this river and the Ohio SQM Project are greatly appreciated. Special thanks are also extended to the Knox Soil and Water Conservation District, Kenyon College Brown Family Environmental Center, private landowners, and the Kokosing Scenic River Advisory Council for their continued support and assistance. These reference stations are also closely monitored by Division of Watercraft staff.

Kokosing River

River Mile 2.5 - Riley Chapel Road Bridge

Bill Abbot

Brian and Keith Robinson

River Mile 8.8 - Bridge Street

Terry Gardner

River Mile 11.6 - Howard Riffle

Jim and Tony Bull

River Mile 16.1 - Zion Road Bridge

Marcia Morgan

Nancy Badet

River Mile 21.3 - Village of Gambier

Joan and Bill Heiser

River Mile 23.2 - Lower Gambier Road

Stephanie, Olivia, and Josie Janosy

River Mile 28.0 - Mt. Vernon Riverside Park

Yelena Arshina

River Mile 32.7 - Beckley Road Bridge

Volunteer Needed

River Mile 40.5 - Lucerne Road Bridge

Erin Carffrey

David Cordray

Kokosing River - North Branch

River Mile 2.1 - Beckley Road Bridge

Not Monitored

River Mile 3.7 - Hyatt Road Bridge

Phyllis and Dave Randall

River Mile 6.3 - Fredericktown Community Park

Loren, Barb and Tyler Ing

Don Schmoeger

The continued success of the Ohio SQM Project is dependent upon the commitment and dedication of these (and past) volunteers. We would like to recognize volunteers *Joan and Bill Heiser; Don Schmoeger; Keith and Brian Robinson; William Abbot; Marcia Morgan; Nancy Badet; Terry Gardner; and Stephanie, Olivia, and Josie Janosy*; for monitoring three times or more during the season. If you are interested in becoming a volunteer, please contact the Central Ohio SQM Coordinator at 740-548-5490 or visit watercraft.ohiodnr.gov/SQM for more information.

Station Descriptions

The Kokosing River and the North Branch of the Kokosing River both largely are surrounded by agriculture. Most land adjacent to the rivers is privately owned and there are few public access sites. When possible, sampling stations are located in or adjacent to areas where public access is permitted. The following are brief descriptions of selected SQM stations on the Kokosing Scenic River.

Kokosing River

River Mile 2.5 - Riley Chapel Road, Butler Township

Located downstream from Riley Chapel Road Bridge in Knox County, this site is the furthest downstream reference station on the Kokosing River. Unlike much of the river, this section lacks a riparian buffer and erosion along the stream banks is apparent. Access to the riffle can be difficult during periods of moderately high water and impossible during periods of high water.

River Mile 8.8 - Bridge Street, Millwood

Upstream from Bridge Street in Millwood, Knox County, this site is accessed on the south side of the bridge. The riffle is wide and a number of different habitats may be sampled with varying depth, flow and substrate composition. However, the current may be very swift in some areas so caution should be exercised.

River Mile 11.6 - Pipesville Road, Howard

This riffle in Knox County is located 200 yards upstream from Pipesville Road. Upon reaching the site, two riffles divided by a gravel bar will be apparent. Monitoring is conducted at the riffle left of the gravel bar as you approach from downstream. Access is via private property with landowner consent.

River Mile 16.10 - Zion Road, Harrison Township

The site is located in Knox County immediately upstream from the Zion Road Bridge. The site is accessible from the eastern side of the bridge, where a path leads to the river. This site has a steep gradient so the water current is quite swift.

River Mile 21.3 – Laymon Road Bridge, Gambier

Running adjacent to the river, the Kokosing Gap Trail is the best way to access this site. 0.30 miles upstream from the Laymon Road Bridge in Knox County the riffle's substrate has a composition of primarily gravel and sand. The smaller substrate makes finding some of the larger macroinvertebrates a challenge; however, the site still yields good biological diversity. Plenty of parking is available at the gravel lot located near the Brown Family Environmental Center.

River Mile 23.2 - Lower Gambier Road, Pleasant Township

Located in Knox County, this riffle is located immediately upstream of Pine Bridge on the Kokosing Gap Trail and continues underneath the bridge. The site is accessed along the Gap Trail or Lower Gambier Rd.

River Mile 28.0 - Riverside Park, Mt. Vernon

Constantly changing due to the Army Corps of Engineers, City of Mount Vernon, and Knox County's dike management. The availability of public access and ample parking space for several vehicles makes this an ideal site for large groups interested in monitoring.

River Mile 32.7 - Beckley Road, Morris Township

Upstream from the Beckley Road Bridge in Knox County, this riffle is located on private property and permission from the landowner is required. River Mile 32.7 was acquired in 2016 to replace a former downstream site, River Mile 32.6. The stream channel was altered significantly at the former site due to bridge maintenance in the fall of 2014. This site was not monitored in 2015 while a new site was being established.

River Mile 40.5 - Lucerne Road, Wayne Township

This riffle is located upstream from the Lucerne Road Bridge in Knox County on private property. There is limited space for one vehicle to park at this site. This site makes up the smallest riffle area (approximately 15 feet by 3 feet) monitored on the Kokosing River. Although small, this site consistently scores in the excellent range.

Kokosing River - North Branch

River Mile 2.1 - Beckley Road, Morris Township

Located upstream from the Beckley Road Bridge in Knox County, this site is on private property and permission from the landowner is necessary. The mouth of this riffle terminates just before a bend in the river and the current can be quite swift at times.

River Mile 3.7 - Hyatt Road, Morris Township

Approximately 100 yards downstream from the Hyatt Road Bridge in Knox County, access is available on the east side of the river along the bridge right-of-way.

River Mile 6.3 - Fredericktown Community Park, Village of Fredericktown

Just downstream from the confluence with the East Branch of the Kokosing River in Knox County, this public site is excellent for large groups and has ample parking. The river is accessed directly from the park.



Sampling Results and General Trends

Average rainfall amounts for the 2015 season produced an average season for monitoring. According to the National Oceanic and Atmospheric Administration (NOAA), during the monitoring months of May through October the central hills region of Ohio saw a total of 22.54" of rain, this is a 1.20" increase over the 100-year average (1916 to 2015) of 21.34". Most of the 2015 samples were taken during the designated sampling periods. The Scenic Rivers SQM Project requires that each assessment be conducted at a minimum of 30 days apart between the months of May and October. Additionally, the SQM Project requires a minimum of three readings to calculate a Cumulative Index Value (CIV). Three sites were not monitored in 2015 for various reasons please see the results below.

During 2015, volunteers and ODNR staff conducted 27 assessments in the Kokosing River. The mainstem of the Kokosing River scored an average CIV of 19.81, corresponding to the good range for stream quality. The CIV average is down from the 2014 average score of 22.93. It is unclear currently why the Kokosing has shown a decline during the past 3 monitoring seasons. Scenic Rivers staff will investigate further into the decline during the 2015 season. Two sites were not monitored on the mainstem in 2015, 11.6 and 32.6, however it is unlikely that they are the source for the decline in the CIV. During the late fall of 2014, the Knox County Engineer removed an island downstream of the Beckley Road bridge. When the island was removed, the riffle was subsequently destroyed. A new monitoring station up stream of the Beckley Road Bridge was discovered during the 2015 season. Permission was granted by the owners of the property in which the riffle is situated. 32.6 will be replaced by 32.7. The Howard monitoring station 11.6 was monitored by a volunteer that was tragically killed during the monitoring season. Unfortunately, by the time the information was discovered it was too late in the season to monitor the site again. The Scenic Rivers' staff sincerely send their condolences to the family and friends of the volunteer lost. The average taxonomic diversity per assessment was 10 macroinvertebrate orders (e.g. stonefly, damselfly, mayfly, etc). The average of taxonomic diversity per assessment has remained consistent for the past three years.

During 2015, volunteers and ODNR staff on North Branch of the Kokosing River conducted 9 assessments. The North Branch scored an average CIV of 24.25, meeting the excellent range of stream quality. This is down slightly from the CIV of 25.56 in 2014. Rivermile 2.1 was not monitored in the 2015. The site had a major substrate shift as a result to a downed tree, we will continue to monitor site over the next two years to see if the area will redevelop a riffle. The North Branch has shown consistent improvement since its designation in 1997. The average taxonomic diversity per assessment was 12 macroinvertebrate orders, up one from the 2014 season average.

Data collected by SQM Project volunteers and ODNR staff is used as a water quality-screening method. The data helps to detect significant changes in stream quality based on CIV data from sites monitored for many years. If there is a significant decline in the average CIV, potential problems that may be causing stream degradation can be investigated and addressed.

Ohio Scenic Rivers Program staff appreciates the assistance received from dedicated volunteer monitors. It is only through their efforts that it was possible to complete the SQM samples on the Kokosing River main stem and North Branch during 2015. Working together has produced significant results but additional Scenic Rivers' volunteers are needed to monitor reference sites, ensuring accurate and thorough data collection. For more information, please contact the Central Ohio SQM Coordinator at 740-548-5490 or find more information at watercraft.ohiodnr.gov/SQM.

Total Suspended Solids (TSS)

In 1999, the Scenic River Program added Total Suspended Solids (TSS) monitoring to the Ohio SQM Project. The purpose of this addition is to estimate the amount of soil sediments affecting a stream by estimating the turbidity of the water. These sediments are attributed to problems originating upstream of the sampling site. The equipment is calibrated to predict TSS at 90% accuracy. The measurements are accurate enough to determine the changes in sediment rates in a stream at a given location and time.

Variables such as amount of precipitation, slope and gradient of the river system, soil type, time of year data is collected, amount of development, amount of riparian corridor, velocity of the river flow, and the amount of waste water effluent have an effect on the TSS value.

Precipitation amount is important because of the increased potential for sediments to be carried into the river during a rain event. The TSS value may appear higher than normal if precipitation amounts are not taken into account. Since large rain events usually happen in the spring and early summer, the time of year the samples are taken could affect the TSS score. The gradient of the stream is important as well. Sediments do not settle out as easily in high gradient streams because the velocity of the water washes it downstream. In low gradient streams, sediment has a chance to settle out, resulting in a lower TSS value. Soil types affect TSS values because some soil types erode faster than others. A better understanding of the types of soils within the watershed may give way to a better understanding of the baseline TSS values for a stream.

Development in an area can cause changes in the TSS score. Areas cleared for new buildings are often not covered, causing an acute rise in the amount of suspended solids in nearby streams. Impermeable surfaces can also cause chronic elevation of TSS values because there is no buffer to absorb or trap runoff. Wastewater treatment plant effluent would only affect TSS scores in low flow situations, and only if the plant employs only primary or secondary treatment.

The actual process of taking a sample is simple. Using a clear Lucite sediment stick developed by the Lake Soil and Water Conservation District, a water sample is collected from the stream. Keeping the sample materials suspended, water is then poured out of the tube until the 0.4-inch target dot is visible on the tube bottom. A reading of the water column height is taken from the markings on the stick to the nearest $\frac{1}{4}$ inch. A conversion table is then used to convert the sediment stick reading to a TSS measurement in the form of an estimate of the weight of solids suspended in the water column (mg/l).

The TSS measurement can be used to estimate water quality with the following scale:

- TSS <10 mg/l = excellent water quality
- TSS 10-28 mg/l = normal water quality
- TSS 29-133 mg/l = impaired water quality
- TSS >133 mg/l = severely impacted water quality

2015 Results: A total of 22 TSS readings were taken in the Kokosing River mainstem. The Kokosing River had a median value of <6.2 mg/L of TSS, corresponding to the maximum stream rating possible for the instrument. The data set ranged from <6.2 to 12 mg/L of total suspended solids. A total of 7 TSS readings were taken in the North Branch of the Kokosing River with a median value of 10 mg/L, corresponding with the normal range. The data set ranged from <6.2 to 66 mg/L.

Comparisons of Collected Stream Quality Monitoring Data

Frequent monitoring of the same reference station is performed a minimum of three times per year consistently year after year. An assessment of the diversity and tolerance levels of taxonomy collected generates the Cumulative Index Value (CIV) for the site on a given date. Field assessment results are used as basic indicators of long-term changes in a stream's macroinvertebrate community and help the Scenic Rivers staff identify pronounced stream quality problems.

Table 1 identifies the 20 macroinvertebrates assessed and their general tolerance to pollutants. Pollution-intolerant organisms, such as those listed in Group I, require unpolluted, high quality water in order to survive. Pollution-tolerant organisms, such as those listed in Group III, are extremely tolerant of deteriorated water conditions.

Table 1. Macroinvertebrate Pollution Tolerance

Group I Taxa Pollution Intolerant	Group II Taxa Moderately Tolerant	Group III Taxa Pollution Tolerant
Water Penny Beetle Larvae (WP) Mayfly Nymphs (MF) Stonefly Nymphs (ST) Dobsonfly Larvae (DO) Caddisfly Larvae (CD) Riffle Beetle Adult (RI) Other Snails (OS)	Damselfly Nymphs (DA) Dragonfly Nymphs (DR) Crane Fly Larvae (CR) Beetle Larvae (BL) Crayfish (CF) Scuds (SC) Clams (CL) Aquatic Sowbugs (SW)	Black Fly Larvae (BF) Aquatic Worms (AW) Midge Larvae (MI) Pouch Snails (PS) Leeches (LE)

Tables 2 and 3 represent the mean CIV for each SQM reference station sampled on the river during 2015. In addition, the table uses symbols (◆) to indicate those macroinvertebrates found to be present at least once during the year at the respective reference station. Each macroinvertebrate is identified by a 2-letter code given in Table 1 (above). A CIV of 23 or greater indicates *Excellent* stream quality; a CIV of 17-22 indicates *Good* stream quality; a CIV of 11-16 suggests *Fair* stream quality; and a CIV of 10 or less reflects *Poor* stream quality. Situated beside the CIVs are the symbols + (improved), = (equal), or – (declined) indicating the relationship to the previous year's CIV.

For the full range of CIVs attained at all sites monitored during the year including non-reference stations, please see the *Appendix*.

Table 2. Kokosing River 2015 Mean CIVs by Reference Station

STATION	W P	M F	S T	D O	C D	R I	O S	D A	D R	C R	B L	C F	S C	S L	S W	B F	A W	M I	P S	L E	CIV
2.5	◆	◆	◆	◆	◆	◆		◆	◆	◆	◆	◆	◆			◆	◆	◆	◆		19+
8.8	◆	◆	◆	◆	◆	◆		◆	◆			◆					◆	◆			17-
11.6																					
16.1	◆	◆	◆		◆	◆	◆		◆	◆	◆	◆	◆	◆		◆	◆	◆	◆	◆	26-
21.3	◆	◆			◆	◆	◆		◆	◆				◆				◆			20-
23.2	◆	◆			◆	◆		◆		◆		◆	◆	◆		◆		◆			17-
28.0	◆	◆	◆		◆	◆	◆		◆	◆	◆	◆	◆	◆		◆	◆	◆	◆		25=
32.6																					
40.5	◆	◆	◆		◆	◆				◆	◆	◆		◆		◆	◆	◆	◆	◆	18-

Table 3. Kokosing River - North Branch 2015 Mean CIVs by Reference Station

STATION	W P	M F	S T	D O	C D	R I	O S	D A	D R	C R	B L	C F	S C	S L	S W	B F	A W	M I	P S	L E	CIV	
2.1																						
3.7	◆	◆	◆	◆	◆	◆	◆			◆	◆	◆	◆	◆		◆	◆	◆				26+
6.3	◆	◆	◆		◆	◆	◆		◆	◆	◆	◆		◆		◆	◆	◆	◆	◆		23-

Figures 1.1 and 1.2 represent the maximum and minimum range of CIVs recorded during the year for each reference station. Figures 2.1 and 2.2 represent mean CIVs at each reference station over many years.

Figure 1.1 - Kokosing River 2015 CIV Maximum and Minimum Ranges




Figure 2.1 - Kokosing River 2006- 2015 Mean CIVs




Figure 1.2 - Kokosing River - North Branch 2014 CIV Maximum and Minimum Ranges





Figure 2.2 - Kokosing River - North Branch 2006-2015 Mean CIVs



Qualitative Habitat Evaluation Index (QHEI)

The Qualitative Habitat Evaluation Index (QHEI) is a system developed and employed by the Ohio Environmental Protection Agency (OEPA) to measure physical habitat conditions in and around rivers and streams in Ohio. During 1998, SQM staff tested a modified version of the QHEI, referred to as *Citizen's QHEI*, to gather baseline measurements at reference stations on several of Ohio's scenic rivers. It is anticipated that such measurements will become yet another annual tool that will be used to monitor habitat and water quality conditions on all Ohio scenic rivers.

Results from 2013 QHEI are included below. When attempting to interpret this data, it is important to recognize that OEPA generally concludes that any site receiving a QHEI value greater than 60 meets current warm water habitat (WWH) standards. Meeting WWH standards suggests that such locations should be adequate for supporting reproducing communities of fish and macroinvertebrate life. Sites attaining QHEI scores of greater than 80 are generally believed to contain exceptional habitat conditions for warm water communities.

The following table has been prepared to assist with determining the relationship between habitat conditions (measured by the QHEI) and macroinvertebrate community performance (measured by the Cumulative Index Value), at each of the reference stations on selected rivers.

Table 4. Kokosing River Mainstem 2013 QHEI and SQM Assessment Data

Reference Station	QHEI	2013 Average CIV	SQM Assessment
RM 2.5	83	17	Good
RM 8.8	91.5	22	Good
RM 11.6	81.5	18	Good
RM 16.1	83.5	28	Excellent
RM 21.3	87	27	Excellent
RM 23.2	93.5	28	Excellent
RM 28.0	83		
RM 32.6	85.5	23	Excellent
RM 40.5	76	25	Excellent

Table 5. Kokosing River- North Branch 2013 QHEI and SQM Assessment Data

Reference Station	QHEI	2013 Average CIV	SQM Assessment
RM 2.1	88	23	Excellent
RM 3.8	66	28	Excellent
RM 6.3	76	29	Excellent

Appendix

2015 Data by Monitoring Station

KOKOSING RIVER - Mainstem																						
RM	DATE	W P	M F	S T	D O	C D	R I	O S	D A	D R	C R	B L	C F	S C	C L	S W	B F	A W	M I	P S	L E	CIV
2.50	5/1/2015		A	A						A							A	A				10
2.50	6/10/2015		C	B	A	B	A							A			C	A	B			20
2.50	6/10/2015		C	B		B	B		A		A	A	A	A			C	A	B			25
2.50	6/10/2015		C	A		B	A							A			C		B			16
2.50	7/23/2015		A		A	A				A	A	A	A	A	A		A		A	A		20
2.50	8/24/2015	A	C	B	A	C	A											B				19
2.50	10/8/2015		A	A		A			A		A	A		A			A		A	A		20
2.50	10/19/2015		A	B	A	C	A				A											17
2.50	10/19/2015		A	B	A	C	A				A											20
8.80	7/7/2015		B			B	A						A					A				12
8.80	8/18/2015	A	C			C	A	A		A			A						B			20
8.80	10/8/2015	A	C	A	A	C			A									A				20
16.10	8/1/2015	B	C			C	B	B		A				A			A	B	B		B	23
16.10	9/7/2015	B	C			B	A	C					A	B	B			B			B	23
16.10	10/22/2015	B	C	A		C	A	A			A	C		B	A		B	B	A	A	A	31
21.30	7/29/2015	B	B			B	A	B		A	A								A			20
21.30	8/30/2015	A	B			C	B	B			B				A				B			20
21.30	10/1/2015	A	B			B	A	A			B				A				A			20
23.20	6/30/2015	A	B			B	A							A			A		B			16
23.20	8/2/2015	A	C			C	B		A		A						B		C			18
23.20	10/18/2015	A	C			C					A		A	B			A					16
28.00	6/24/2015	A	A	A		C	A	A			A	A					A		A			24
28.00	8/26/2015	A	B			C	B	A				B	A		A		B	A	B			24
28.00	10/3/2015	A	A			A	A	B		A			A	A	A		A		A	A		26
40.50	7/19/2015	A		A			B				A								A	A	A	14
40.50	9/7/2015	A	A			C	C					B	C	A				A		A		20
40.50	11/4/2015	A	B	A		C	A					B	A				A	B	B			21

KOKOSING RIVER -NORTH BRANCH																						
RM	DATE	W P	M F	S T	D O	C D	R I	O S	D A	D R	C R	B L	C F	S C	C L	S W	B F	A W	M I	P S	L E	CIV
3.70	9/19/2015	A		A	A	A		A			A		A	A	A		A	A	A			26
3.70	8/18/2015	B	B	A		B	A	A				B	A		A		A	A	A			27
3.70	10/20/2015	A		A		A	A	A			A		A	A	A		A	A				25
6.30	7/22/2015	B	B			C	B	A				B	A		B		B	B	B	A		25
6.30	7/11/2015	B	A			C	B			A	A	B	B		A		B	B	B		A	26
6.30	9/1/2015	C	A	A		C	A				A	B			A		A	B	B			24
6.30	9/26/2015	B	B				B	A				B	A		B		A	B	B	A		22
6.30	10/8/2015	B	B	A		B						B			A		A	B	B			19